

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence selected from the group consisting of:
 - (a) a polynucleotide fragment of SEQ ID NO:1 or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No: PTA-4454, which is hybridizable to SEQ ID NO:1;
 - (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:2 or a polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No: PTA-4454, which is hybridizable to SEQ ID NO:1;
 - (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:2 or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No: PTA-4454, which is hybridizable to SEQ ID NO:1;
 - (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:2 or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No: PTA-4454, which is hybridizable to SEQ ID NO:1;
 - (e) a polynucleotide encoding a polypeptide of SEQ ID NO:2 or the cDNA sequence included in ATCC Deposit No: PTA-4454, which is hybridizable to SEQ ID NO:1, having tyrosine tubulin ligase activity;
 - (f) a polynucleotide which is a variant of SEQ ID NO:1;
 - (g) a polynucleotide which is an allelic variant of SEQ ID NO:1;
 - (h) an isolated polynucleotide comprising nucleotides 156 to 1775 of SEQ ID NO:1, wherein said nucleotides encode a polypeptide corresponding to amino acids 2 to 541 of SEQ ID NO:2 minus the start methionine;
 - (i) an isolated polynucleotide comprising nucleotides 153 to 1775 of SEQ ID NO:1, wherein said nucleotides encode a polypeptide corresponding to amino acids 2 to 541 of SEQ ID NO:2 including the start codon;
 - (j) an isolated polynucleotide comprising nucleotides 369 to 1247 of SEQ ID NO:1, wherein said nucleotides encode a polypeptide corresponding to amino acids 73 to 365 of SEQ ID NO:2 including the start codon;
 - (k) an isolated polynucleotide comprising a polynucleotide that encodes at least 424 contiguous amino acids of SEQ ID NO:2;

(l) an isolated polynucleotide comprising at least 1272 contiguous nucleotides of SEQ ID NO:1;

(m) a polynucleotide which represents the complimentary sequence (antisense) of SEQ ID NO:1;

5 (n) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(m), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues;

(o) a polynucleotide comprising the BGS-42 gene;

10 (p) a polynucleotide comprising the BGS-42 promoter; and

(q) a polynucleotide comprising the nucleotide sequence provided as SEQ ID NO:27.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment consists of a nucleotide sequence encoding a human tyrosine
15 tubulin ligase.

3. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.

4. A recombinant host cell comprising the vector sequences of claim 3.

5. An isolated polypeptide comprising an amino acid sequence selected
20 from the group consisting of:

(a) a polypeptide fragment of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-4454;

(b) a polypeptide fragment of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-4454, having tyrosine tubulin ligase activity;

25 (c) a polypeptide domain of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-4454;

(d) a polypeptide epitope of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-4454;

30 (e) a full length protein of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: PTA-4454;

(f) a polypeptide comprising amino acids 2 to 541 of SEQ ID NO:2, wherein said amino acids 2 to 541 comprising a polypeptide of SEQ ID NO:2 minus the start methionine;

(g) a polypeptide comprising amino acids 1 to 541 of SEQ ID NO:2;

5 (h) a polypeptide comprising amino acids 73 to 365 of SEQ ID NO:2; and

(i) a polypeptide comprising at least 424 contiguous amino acids of SEQ ID NO:2.

6. The isolated polypeptide of claim 5, wherein the full length protein comprises sequential amino acid deletions from either the C-terminus or the N-terminus.

7. An isolated antibody that binds specifically to the isolated polypeptide of claim 5.

8. A recombinant host cell that expresses the isolated polypeptide of claim 5.

15 9. A method of making an isolated polypeptide comprising:

(a) culturing the recombinant host cell of claim 8 under conditions such that said polypeptide is expressed; and

(b) recovering said polypeptide.

10. The polypeptide produced by claim 9.

20 11. A method for preventing, treating, or ameliorating a medical condition, comprising the step of administering to a mammalian subject a therapeutically effective amount of the polypeptide of claim 5, or a modulator thereof.

12. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

25 (a) determining the presence or absence of a mutation in the polynucleotide of claim 1; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

30 13. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the polypeptide of claim 5 in a tyrosine tubulin ligase sample; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

14. An isolated nucleic acid molecule consisting of a polynucleotide having a nucleotide sequence selected from the group consisting of:

- 5 (a) a polynucleotide encoding a polypeptide of SEQ ID NO:2;
- (b) an isolated polynucleotide consisting of nucleotides 156 to 1775 of SEQ ID NO:1, wherein said nucleotides encode a polypeptide corresponding to amino acids 2 to 541 of SEQ ID NO:2 minus the start codon;
- (c) an isolated polynucleotide consisting of nucleotides 369 to 1247 of SEQ ID NO:1, wherein said nucleotides encode a polypeptide corresponding to amino acids 73 to 365 of SEQ ID NO:2 including the start codon;
- (d) a polynucleotide encoding the BGS-42 polypeptide encoded by the cDNA clone contained in ATCC Deposit No. PTA-4454;
- (e) a polynucleotide which represents the complimentary sequence (antisense) of SEQ ID NO:1;
- 15 (f) an isolated polynucleotide consisting of a polynucleotide that encodes at least 424 contiguous amino acids of SEQ ID NO:2;
- (g) an isolated polynucleotide consisting of at least 1272 contiguous nucleotides of SEQ ID NO:1;
- 20 (h) a polynucleotide consisting of the BGS-42 gene;
- (i) a polynucleotide consisting of the BGS-42 promoter; and
- (j) a polynucleotide comprising the nucleotide sequence provided as SEQ ID NO:27.

25 15. The isolated nucleic acid molecule of claim 14, wherein the polynucleotide comprises a nucleotide sequence encoding a human tyrosine tubulin ligase.

 16. A recombinant vector comprising the isolated nucleic acid molecule of claim 15.

 17. A recombinant host cell comprising the recombinant vector of claim 16.

30 18. An isolated polypeptide consisting of an amino acid sequence selected from the group consisting of:

(a) a polypeptide fragment of SEQ ID NO:2 having tyrosine tubulin ligase activity;

(b) a polypeptide domain of SEQ ID NO:2 having tyrosine tubulin ligase activity;

5 (c) a full length protein of SEQ ID NO:2;

(d) a polypeptide corresponding to amino acids 2 to 541 of SEQ ID NO:2, wherein said amino acids 2 to 541 consisting of a polypeptide of SEQ ID NO:2 minus the start methionine;

(e) a polypeptide corresponding to amino acids 1 to 541 of SEQ ID NO:2;

10 (f) a polypeptide corresponding to amino acids 73 to 365 of SEQ ID NO:2; and

(g) a polypeptide encoded by the cDNA contained in ATCC Deposit No. PTA-4454.

15 19. The method of diagnosing a pathological condition of claim 15 wherein the condition is a member of the group consisting of: a disorder related to aberrant tubulin ligase activity; a disorder related to aberrant tubulin-carboxypeptidase activity; aberrant cellular proliferation; reproductive disorders; testicular disorders; testicular cancer; pulmonary disorders; lung cancer; gastrointestinal disorders; colon cancer; stomach cancer; neural disorders; brain cancer; liver cancer; proliferative condition of the testis; proliferative condition of the lung; proliferative condition of the small intestine; proliferative condition of the brain; and proliferative condition of the lymph tissue.

20 20. The method for preventing, treating, or ameliorating a medical condition of claim 11, wherein the medical condition is selected from the group consisting of: a disorder related to aberrant tubulin ligase activity; a disorder related to aberrant tubulin-carboxypeptidase activity; aberrant cellular proliferation; reproductive disorders; testicular disorders; testicular cancer; pulmonary disorders; lung cancer; gastrointestinal disorders; colon cancer; stomach cancer; neural disorders; brain cancer; liver cancer; proliferative condition of the testis; proliferative condition of the lung; proliferative condition of the small intestine; proliferative condition of the brain; and proliferative condition of the lymph tissue.

25 30

21. A computer for producing a three-dimensional representation of a molecule or molecular complex, wherein said molecule or molecular complex comprises the structural coordinates of the BGS-42 model provided in Figure 13 in accordance with Table IV wherein said computer comprises:

5 (a) A machine-readable data storage medium, comprising a data storage material encoded with machine readable data, wherein the data is defined by the set of structure coordinates of the model;

 (b) a working memory for storing instructions for processing said machine-readable data;

10 (c) a central-processing unit coupled to said working memory and to said machine-readable data storage medium for processing said machine readable data into said three-dimensional representation; and

 (d) a display coupled to said central-processing unit for displaying said three-dimensional representation.

15 22. A method for identifying a mutant with altered biological properties, function, or activity of BGS-42 wherein said method comprises the steps of:

 (a) using a model of said polypeptide according to the structural coordinates of said model to identify amino acids to mutate; and

20 (b) mutating said amino acids to create a mutant protein with altered biological function or properties.

23. A method for designing or selecting compounds as potential modulators of BGS-42 wherein said method comprises the steps of:

 (a) identifying a structural or chemical feature of said member using the structural coordinates of said member; and

25 (b) rationally designing compounds that bind to said feature.